

Browser:
IP Address:
Unique ID:
Location:

Chrome 64.0.3282.140 / Windows
40.439800262451, -86.089302062988

You are not alone. We are here to help.

Are you interested in learning more about clinical trials and treatment options that might be available to the patient based on your answers to this questionnaire? Yes

If the patient elects to be in a research study do they have a spouse, caregiver, or friend who lives with them that could help with study participation and could attend some study visits? Yes

What best describes you? I am the patient

The patient has been diagnosed with: Parkinson's Disease

This information will help identify opportunities that might best match the patient's disease progression.

How many years ago was the patient diagnosed with Parkinson's disease? 6 years ago

What year was the patient diagnosed with Parkinson's Disease? 2012

Contact Information

Patient Name [REDACTED]

Age 54

Address [REDACTED] IN 46902

Email [REDACTED]@aol.com

Phone (765) [REDACTED]

Who is the patient's current neurologist? Dr. [REDACTED]

Agreement I have read and agree to the Parkinsons.Community Privacy Policy and Terms of Use

OFF Time Questions

Can the patient tell when the levodopa/carbidopa wears off? Yes

Does OFF time interfere with day to day activities or quality of life? Yes

Is OFF time a problem for the patient? Yes

During OFF times, would the patient be willing to consider using a pen device to get moving again? Yes

Is the patient scheduled for Deep Brain Stimulation (DBS) within the next 3 months? No

Important Medication Questions

Is the patient taking levodopa (i.e. Sinemet, Rytary, etc)? Yes, the patient has taken levodopa / carbidopa for MORE than 30 days.

How many times/day is the patient taking Sinemet or Rytary? Sinemet 4 times or more
Both Rytary and Sinemet

Is the patient taking or have they previously taken the following medications? Dopamine Agonists (Mirapex/ Pramipexole. Requip/ Ropinirole, Neupro Patch/ Rotigotine) = Currently Taking
COMT Inhibitors (Comtan/ Stalevo/ Entacapone) Usually an orange pill. = Previously Taken and Stopped
Amantadine (including Gocovri) = Currently Taking

Is the patient taking or have they previously taken Apokyn? Never Taken

Does the patient have a Duopa pump? No

These questions are to help determine the basic eligibility status of the patient.

Is the patient currently participating in a clinical trial? No

Does the patient have a history of frequent nausea or vomiting? No

Has the patient had difficulties tolerating any PD medication due to frequent nausea or vomiting in the past? No

Has the patient been diagnosed with dementia or have significant trouble with memory or thinking that affects day to day activities (i.e. cognitive impairment)

None

Within the past two years has the patient taken antipsychotic medication? (e.g. Seroquel/ quetiapine, Nuplazid/ pimavanserin, Clozaril/ clozapine, Risperdal/ risperidone, Zyprexa/ olanzapine)?

No

Does the patient have either of the following psychiatric conditions?

Neither

Does the patient currently have or in the past had hallucinations (e.g. seeing people, animals, or things are not truly present) or delusions (fixed unusual thoughts such as paranoia)?

No

You are doing great.

Has the patient had brain surgery?

No

Has the patient ever had a stroke?

No

Has the patient ever had any type of cancer, even skin cancer?

Yes

What type of cancer?

basil carcinoma

When was the patient diagnosed with cancer?

2012

Did this cancer occur in the past 5 years?

No

Does the patient have kidney dialysis or has their doctor diagnosed them with any acute or chronic renal insufficiency or failure?

No

Does the patient have a history of recent significant cardiac problems: heart attack within the last year, bypass surgery in the last year, pacemaker placement in the past year?

No

This is the last set of general questions.

Is the patient currently using any cannabis products (Medical or Recreational) by smoking/ vaping. Edibles, or topicals to skin? Yes, but would be willing to stop using these products to participate in a clinical trial with and investigational medication for PD

Has the patient been diagnosed with addiction or treated for use of any addictive drug (e.g, opioids/ narcotics. methamphetamine, cocaine, etc,)? No

Has the patient been diagnosed with or treated for excessive alcohol use (alcoholism) in the last 2 years? No

Have you ever been diagnosed with impulse control disorder associated with use of PD medication (especially dopamine agonists such as Mirapex/ pramipexole, Requip/ ropinirole, or Neupro patch/ rotigotine patch)? No

Do you or have you ever felt driven to do or think about something and find it hard to stop such as gambling, cleaning, using the computer, taking extra medicine, obsessing about food or sex? No

Does the patient have bothersome drooling? No

Would the patient like to consider an investigational medication for treatment symptoms of lightheadedness or orthostatic hypotension? Yes

Does the patient have symptoms of lightheadedness or feeling like they might faint frequently when standing up from a chair or lying? Yes

Has the patient been diagnosed as having orthostatic hypotension (excessive blood pressure drop when you stand up from lying or sitting)? No

Has the patient been diagnosed with sleep apnea? No

Is the patient currently or have they been treated with any of the following medications for orthostatic hypotension?

Midodine/ Proamatine = No
Florinef/ fludrocortisone = No
Northera/ droxidopa = No
Mestinon/ pyridostigmine = No
Strattera/ atomoxetine = No

Is the patient currently or have they been treated with any of the following non-drug therapies for orthostatic hypotension?

Extra fluid intake = No
Extra salt intake = No
Elevation of head of bed = No
Compression stockings = No
Extra caffeine intake = No

Has the patient fainted (lost consciousness and fell down) as a result of feeling lightheaded when standing up from a chair or lying down since diagnosed with Parkinson's disease and these problems began?

No

Sleep Related Questions

Would the patient like to consider an investigational medication for the treatment of symptoms related to sleepiness and Parkinson's disease?

Yes

Please rate the chance of dozing off in the following situations over the past few weeks:

Sitting and reading = 3 high chance of dozing
Watching TV = 3 high chance of dozing
Sitting inactive in a public place (e.g a theater or a meeting) = 2 moderate chance of dozing
As a passenger in a car for an hour without a break = 2 moderate chance of dozing
Lying down to rest in the afternoon when circumstances permit = 2 moderate chance of dozing
Sitting and talking to someone = 1 slight chance of dozing
Sitting quietly after a lunch without alcohol = 2 moderate chance of dozing
In a car, while stopped for a few minutes in traffic = 2 moderate chance of dozing

Based on your answers you have indicated that the patient likely meets the criteria to speak with a Parkinson's Community Advocate. Would you like to see the available resources?

Yes, let's continue.
